

Cranial nerves

Organize first. Then test with purpose.

A first-pass study guide for PT, PTA, and DPT learners. Use it to connect function, broad region, and a focused observation. Do not use one finding to diagnose a person.

THE LEARNING LOOP

SORT	>	MAP	>	APPLY	>	RETRIEVE	>	CHECK
function		region		observation		without notes		reasoning

AFTER REVIEWING THIS GUIDE, YOU SHOULD BE ABLE TO

- 1 Sort**
Classify the 12 nerves using the traditional sensory, motor, and mixed framework.
- 2 Map**
Place the main brainstem groups in a simplified midbrain, pons-region, medulla sequence.
- 3 Apply**
Connect a focused observation to a nerve group, then state why context still matters.


HIGH-YIELD FIRST PASS

Ask what system is affected. Group the nerve by function and broad region. Then pause: a pattern can guide a focused exam, but it does not establish a diagnosis.

Use the next page as your 30-second route. Use Page 3 to practice the pause.

See the route in 30 seconds

A simplified learning map, not a complete nuclei or lesion map.



Fast cue: 3-4 midbrain / 5-8 pons region / 9-12 medulla

SORT BEFORE DETAILS

SENSORY

I II VIII

MOTOR

III IV VI XI XII

MIXED

V VII IX X

Traditional categories help you organize first. Individual sensory, motor, and autonomic components still matter.

A focused observation needs a calm sequence

Use the sequence to organize a study prompt. The final step protects against overcalling.

FOUR STEPS

- 1 Name the system**
What is the prompt asking you to observe?
- 2 Group likely nerves**
Use function before a memorized detail.
- 3 Locate broadly**
Retrieve the route cue, not a diagnosis.
- 4 Pause for context**
Pattern, timing, associated signs, and the full exam change meaning.

THREE PRACTICE STARTS

Lateral eye movement

Start with III-IV-VI. Then retrieve: VI supplies lateral rectus.

PAUSE: A single ocular finding does not assign one lesion.

Facial sensation + mastication

Start with V, a mixed nerve. Add its three sensory divisions during course review.

PAUSE: Pattern recognition is not a diagnosis.

Shoulder elevation + head turn

Start with XI, a motor nerve. Connect sternocleidomastoid and trapezius.

PAUSE: Interpret with the rest of the neurologic exam.

FOCUSED OBSERVATION INDEX

I-II Smell / vision: use relevant history or visual function in the appropriate exam context.

III-IV-VI Extraocular movements and pupil-related findings belong together.

V-VII Compare facial sensation or movement thoughtfully, with symmetry in view.

VIII Connect hearing or vestibular observations to the exam question.

IX-X Voice, palate, and swallow-related features require supervision and setting context.

XI-XII Connect neck/shoulder or tongue movement to the focused examination.

DO NOT OVERCALL: a study observation identifies a question for the focused exam. It does not diagnose a person or make a swallow, airway, or management decision.

Retrieve first. Then check your reasoning.

Cover the lower half before you answer. Write a cue, not a paragraph.

1. Which three cranial nerves are traditionally sensory only?

2. What is the fastest broad-region sequence for III-XII?

3. A learner notices facial movement asymmetry. What is the safe next thought?

CHECK

1. I, II, VIII.
2. III-IV midbrain; V-VIII pons region; IX-XII medulla.
3. Connect the observation to the facial-movement group and examine the pattern in context. One finding does not establish a diagnosis.

IF YOU MISSED THIS

Return to Page 2 for the route and sort. Return to Page 3 for the pause/context step. Then retrieve again before reopening the answer key.

SOURCES AND SCOPE

1. Sonne J, Omole AE, Lopez-Ojeda W. Neuroanatomy, Cranial Nerve. StatPearls / NCBI Bookshelf.
2. Rohde S, et al. Evaluation of Patients with Cranial Nerve Disorders. NCBI Bookshelf.
3. Yaddanapudi S. How To Do the Cranial Nerve Examination. Merck Manual Professional Edition.

Educational anatomy and reasoning only. This does not replace course instruction, supervision, an examination protocol, medical advice, or clinical judgment.